

COMPACT HANKEL OPERATORS ON GENERALIZED BERGMAN SPACES OF THE POLYDISC

TRIEU LE

ABSTRACT. Let ϑ be a measure on the polydisc $\mathbb{D}^n$ which is the product of n regular Borel probability measures so that $\vartheta([r, 1]^n \times \mathbb{T}^n) > 0$ for all $0 < r < 1$. The Bergman space A_{ϑ}^2 consists of all holomorphic functions that are square integrable with respect to ϑ . In one dimension, it is well known that if f is continuous on the closed disc $\overline{\mathbb{D}}$, then the Hankel operator H_f is compact on A_{ϑ}^2 . In this paper we show that for $n \geq 2$ and f a continuous function on $\overline{\mathbb{D}^n}$, H_f is compact on A_{ϑ}^2 if and only if there is a decomposition $f = h + g$, where h belongs to A_{ϑ}^2 and $\lim_{z \rightarrow \partial \mathbb{D}^n} g(z) = 0$.

1. INTRODUCTION

Fix a positive integer $n \geq 1$. Let $\mathbb{D}^n$ be the open unit polydisc in $\mathbb{C}^n$ and let $\mathbb{T}^n$ be the n -torus, which is the Shilov boundary of $\mathbb{D}^n$. The closure of $\mathbb{D}^n$ is $\overline{\mathbb{D}^n}$, the product of n copies of the closed unit disc.

For $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ and $\zeta = (\zeta_1, \dots, \zeta_n) \in \mathbb{T}^n$, we use $z \cdot \zeta$ and $\zeta \cdot z$ to denote the point $(z_1 \zeta_1, \dots, z_n \zeta_n)$. We write $\bar{z} = (\bar{z}_1, \dots, \bar{z}_n)$, and for any $m = (m_1, \dots, m_n)$ in $\mathbb{Z}^n$, we write $z^m = z_1^{m_1} \dots z_n^{m_n}$ whenever it is defined. We use σ to denote the surface measure on $\mathbb{T}^n$ which is normalized so that $\sigma(\mathbb{T}^n) = 1$. Let μ be a regular Borel probability measure on $[0, 1]^n$. Then there is a regular Borel probability measure ϑ on $\mathbb{D}^n$ so that

$$\int_{\mathbb{D}^n} f(z) d\vartheta(z) = \int_{[0,1]^n} \left\{ \int_{\mathbb{T}^n} f(r \cdot \zeta) d\sigma(\zeta) \right\} d\mu(r) \quad (1.1)$$

for all continuous functions f with compact support on $\mathbb{D}^n$. It then follows that the above identity also holds true for all functions f in $L^1(\mathbb{D}^n, \vartheta)$.

In this paper we are interested in those measures μ which satisfy the condition $\mu([r, 1]^n) > 0$ for $0 < r < 1$. This implies that $\vartheta(\{z \in \mathbb{D}^n : |z_1| \geq r, \dots, |z_n| \geq r\}) > 0$ for $0 < r < 1$. We write L_{ϑ}^2 for $L^2(\mathbb{D}^n, \vartheta)$ and $\|\cdot\|_2$ for the norm on L_{ϑ}^2 . The Bergman space A_{ϑ}^2 is the closure in L_{ϑ}^2 of the space of all holomorphic polynomials. The condition on μ will imply that all functions in A_{ϑ}^2 are holomorphic on the polydisc. Let P denote the orthogonal projection from L_{ϑ}^2 onto A_{ϑ}^2 . For any function f in L_{ϑ}^2 , the (big) Hankel operator H_f is a densely defined operator from A_{ϑ}^2 into $L_{\vartheta}^2 \ominus A_{\vartheta}^2$ by $H_f(\varphi) = (1 - P)(f\varphi)$ for all holomorphic polynomials φ . The function f

1991 *Mathematics Subject Classification.* Primary 47B35.

Key words and phrases. Bergman space, Hankel operator, compactness.

will be called a symbol of the operator H_f . It is clear that if f is bounded, then H_f is a bounded operator with $\|H_f\| \leq \|f\|_\infty$. However, there are unbounded functions f such that H_f extends to a bounded operator on A_ϑ^2 . In fact, if f belongs to A_ϑ^2 , then since $f\varphi$ belongs to A_ϑ^2 for all holomorphic polynomials φ , we conclude that $H_f = 0$. Conversely, if $H_f = 0$, then since $0 = H_f(1) = (I - P)(f)$, we see that f must belong to A_ϑ^2 . Therefore, $H_f = 0$ if and only if f is in A_ϑ^2 . This shows that a Hankel operator has many symbols and any two symbols of the same operator differ by a function in A_ϑ^2 .

It is well known that if a function $g \in L_\vartheta^2$ vanishes outside a compact subset of $\mathbb{D}^n$, then the Hankel operator H_g is compact. Let $\partial\mathbb{D}^n$ be the topological boundary of $\mathbb{D}^n$ as a subset of $\mathbb{C}^n$. If $g \in L_\vartheta^2$ such that $\lim_{z \rightarrow \partial\mathbb{D}^n} g(z) = 0$ (that is, for any $\epsilon > 0$, there is a compact subset M of $\mathbb{D}^n$ so that $|g(z)| < \epsilon$ for ϑ -a.e. z in $\mathbb{D}^n \setminus M$), then an approximation argument shows that H_g is also a compact operator. This together with the above fact about zero Hankel operators implies that if $f = h + g$, where h belongs to A_ϑ^2 and $\lim_{z \rightarrow \partial\mathbb{D}^n} g(z) = 0$, then H_f is compact.

In the one-dimensional case ($n = 1$), it is well known that if f is continuous on $\overline{\mathbb{D}}$, then H_f is compact. See [11, p. 226] for the case of weighted Bergman spaces. For generalized Bergman spaces, one can prove this by checking directly that $H_{\bar{z}^i z^j}$ is compact for all integers $i, j \geq 0$. See Section 3 for more details. The case $n \geq 2$ turns out to be completely different. Not all Hankel operators with continuous symbols are compact. More surprisingly, we will show, under the assumption that μ is the product of n measures on $[0, 1]$, that if f is continuous on $\overline{\mathbb{D}^n}$, then H_f is compact if and only if f admits a decomposition $f = h + g$, where h belongs to A_ϑ^2 and $\lim_{z \rightarrow \partial\mathbb{D}^n} g(z) = 0$.

If $d\mu(r_1, \dots, r_n) = 2^n r_1 \cdots r_n dr_1 \cdots dr_n$, then A_ϑ^2 is the usual Bergman space of the polydisc. K. Stroethoff [8, 9] and D. Zheng [10] gave necessary and sufficient conditions on a bounded function f for which H_f is compact. However, their conditions, which involve the projection P and Mobius transformations, are difficult to check. Indeed, even if a function f is assumed to be continuous on $\overline{\mathbb{D}^n}$, it is not clear from their results what geometric conditions f needs to satisfy in order for H_f to be compact. Our approach (though works only for continuous functions) is different from theirs and our result is more transparent.

To conclude the section, we would like to mention some results on the compactness of Hankel operators on the Hardy space $H^2 = H^2(\mathbb{T}^n)$. In the one-dimensional case, it is a classical theorem of Hartman (see [11, Chapter 10]) that H_f can be extended to a compact operator if and only if $f = h + g$, where h belongs to H^2 and g is continuous on the circle $\mathbb{T}$. On the other hand, the case $n \geq 2$ is much different. It was showed by M. Cotlar and C. Sadosky in [4] and P. Ahern, E.H. Youssfi and K. Zhu in [1] with a different approach that if H_f is compact, then f must belong to H^2 . This means that there is no non-zero compact Hankel operator on H^2 . This result was also

proved in the setting of Hardy-Sobolev spaces on the polydisc by Ahern, Youssfi and Zhu in the same paper with the same approach. Our analysis in the present paper was actually motivated by their results and techniques.

2. PRELIMINARIES

In this section we explain in more details some of the results that we mentioned in the Introduction. From Cauchy's formula and the assumption that $\mu([r, 1)^n) > 0$ for all $0 < r < 1$, for any compact subset M of $\mathbb{D}^n$, there is a positive constant C_M so that $|p(z)| \leq C_M \|p\|_2$ for all $z \in M$, and all holomorphic polynomials p . This implies that for $f \in A_\vartheta^2$, f is holomorphic on $\mathbb{D}^n$ and we also have $|f(z)| \leq C_M \|f\|_2$ for all $z \in M$. In fact, it can be showed that A_ϑ^2 is the space of all functions in L_ϑ^2 that are holomorphic on $\mathbb{D}^n$. Since $|f(z)| \leq C_M \|f\|_2$, the valuation map $z \mapsto f(z)$ is a continuous functional on A_ϑ^2 for any $z \in \mathbb{D}^n$. So there is a function K_z in A_ϑ^2 such that $f(z) = \langle f, K_z \rangle$ for any $f \in A_\vartheta^2$. The function K_z is called the reproducing kernel at z . For any compact subset M and for any $z \in M$, since $K_z(z) \leq C_M \|K_z\|_2 = C_M (K_z(z))^{1/2}$, we have $K_z(z) \leq C_M^2$.

From (1.1), the monomials $\{z^m : m \in \mathbb{Z}_+^n\}$ are pairwise orthogonal. On the other hand, the linear span of these monomials is dense in A_ϑ^2 . Therefore A_ϑ^2 has the following orthonormal basis, usually referred to as the standard orthonormal basis, $\{e_m(z) = \frac{z^m}{\sqrt{c_m}} : m \in \mathbb{Z}_+^n\}$, where

$$c_m = \int_{\mathbb{D}^n} z^m \bar{z}^m d\vartheta(z) = \int_{[0,1]^n} r_1^{2m_1} \cdots r_n^{2m_n} d\mu(r_1, \dots, r_n).$$

Suppose f is a function in L_ϑ^2 . Then

$$\begin{aligned} \sum_{m \in \mathbb{Z}_+^n} \|H_f e_m\|_2^2 &\leq \sum_{m \in \mathbb{Z}_+^n} \|f e_m\|_2^2 \\ &= \int_{\mathbb{D}^n} |f(z)|^2 \left\{ \sum_{m \in \mathbb{Z}_+^n} |e_m(z)|^2 \right\} d\vartheta(z) \\ &= \int_{\mathbb{D}^n} |f(z)|^2 K_z(z) d\vartheta(z), \end{aligned} \tag{2.1}$$

where the last equality follows from the well known formula

$$K_z(z) = \|K_z\|_2^2 = \sum_{m \in \mathbb{Z}_+^n} |\langle K_z, e_m \rangle|^2 = \sum_{m \in \mathbb{Z}_+^n} |e_m(z)|^2.$$

If f vanishes outside a compact subset M of $\mathbb{D}^n$, then (2.1) gives

$$\sum_{m \in \mathbb{Z}_+^n} \|H_f e_m\|_2^2 \leq C_M^2 \int_M |f(z)|^2 d\vartheta(z) < \infty,$$

since $K_z(z) \leq C_M^2 < \infty$ for all $z \in M$. Thus, H_f is a Hilbert-Schmidt operator, hence it is compact.

Suppose f belongs to L^2_{ϑ} so that $\lim_{z \rightarrow \partial \mathbb{D}^n} f(z) = 0$. Then for any $\epsilon > 0$, there is a compact subset $M_\epsilon \subset \mathbb{D}^n$ so that $|f(z)| < \epsilon$ for ϑ -a.e. $z \in \mathbb{D}^n \setminus M_\epsilon$. This implies that $\|f - f\chi_{M_\epsilon}\|_\infty \leq \epsilon$. And hence, $\|H_f - H_{f\chi_{M_\epsilon}}\| \leq \epsilon$. As we have seen above, $H_{f\chi_{M_\epsilon}}$ is a compact operator for each ϵ . Therefore, H_f , being the limit of a net of compact operators, is also a compact operator. Thus we have showed the following well known result.

Proposition 2.1. *Suppose $f = h + g$, where $h \in A^2_{\vartheta}$ and $g \in L^2_{\vartheta}$ so that $\lim_{z \rightarrow \partial \mathbb{D}^n} g(z) = 0$. Then H_f is a compact operator on A^2_{ϑ} .*

In the rest of the section, we study a decomposition of L^2_{ϑ} into pairwise orthogonal subspaces. If a function belongs to one of these subspaces, the corresponding Hankel operator has a simple form which we can analyze easily. This is one of the key points in our study of the compactness of Hankel operators with continuous symbols.

For any n -tuple $l \in \mathbb{Z}^n$, let $\mathcal{H}_l$ be the space of all functions f in L^2_{ϑ} such that for all $\zeta \in \mathbb{T}^n$, $f(\zeta \cdot z) = \zeta^l f(z)$ for ϑ -a.e. $z \in \mathbb{D}^n$. Following [7], we call each function in $\mathcal{H}_l$ quasi-homogeneous of multi-degree l . It is clear that $\mathcal{H}_l$ is a closed subspace of L^2_{ϑ} . Let Q_l denote the orthogonal projection from L^2_{ϑ} onto $\mathcal{H}_l$. The following lemma shows that these projections are pairwise orthogonal and they constitute a partition of the identity.

Lemma 2.2. *For $s \in \mathbb{Z}^n$ and $f \in L^2_{\vartheta}$, we have*

$$(Q_s f)(z) = \int_{\mathbb{T}^n} f(z \cdot \zeta) \bar{\zeta}^s d\sigma(\zeta), \quad (2.2)$$

for ϑ -a.e. $z \in \mathbb{D}^n$. Furthermore, $\mathcal{H}_l \perp \mathcal{H}_s$ (which implies $Q_l Q_s = 0$) whenever $l \neq s$, and $L^2_{\vartheta} = \bigoplus_{s \in \mathbb{Z}^n} Q_s(L^2_{\vartheta}) = \bigoplus_{s \in \mathbb{Z}^n} \mathcal{H}_s$.

Proof. Since f belongs to L^2_{ϑ} , the integral on the right hand side of (2.2) is well-defined for ϑ -a.e. $z \in \mathbb{D}^n$. For such z , let $f_s(z)$ be the value of the integral. For other values of z , let $f_s(z) = 0$. We will show $Q_s f = f_s$ by proving that $f_s \in \mathcal{H}_s$ and $(f - f_s) \perp \mathcal{H}_s$. For z and any $\gamma \in \mathbb{T}^n$, if the integral in (2.2) is defined, by the rotation invariance of σ , we have

$$\begin{aligned} f_s(z \cdot \gamma) &= \int_{\mathbb{T}^n} f((z \cdot \gamma) \cdot \zeta) \bar{\zeta}^s d\sigma(\zeta) = \int_{\mathbb{T}^n} f(z \cdot (\gamma \cdot \zeta)) \bar{\zeta}^s d\sigma(\zeta) \\ &= \int_{\mathbb{T}^n} f(z \cdot \zeta) \gamma^s d\sigma(\zeta) = \gamma^s f_s(z). \end{aligned}$$

If the integral in (2.2) is not defined, then $f_s(z \cdot \gamma) = \gamma^s f_s(z)$ because they are both zero. Therefore, f_s belongs to $\mathcal{H}_s$.

Now suppose g is a function in $\mathcal{H}_s$. Then

$$\begin{aligned}
\int_{\mathbb{D}^n} f_s(z) \bar{g}(z) d\vartheta(z) &= \int_{\mathbb{D}^n} \int_{\mathbb{T}^n} f(z \cdot \zeta) \bar{\zeta}^s \bar{g}(z) d\sigma(\zeta) d\vartheta(z) \\
&= \int_{\mathbb{T}^n} \int_{\mathbb{D}^n} f(z \cdot \zeta) \bar{\zeta}^s \bar{g}(z) d\vartheta(z) d\sigma(\zeta) \\
&= \int_{\mathbb{T}^n} \int_{\mathbb{D}^n} f(z \cdot \zeta) \bar{g}(z \cdot \zeta) d\vartheta(z) d\sigma(\zeta) \\
&\quad (\text{since } g(z \cdot \zeta) = \zeta^s g(z) \text{ for } \vartheta\text{-a.e. } z) \\
&= \int_{\mathbb{T}^n} \int_{\mathbb{D}^n} f(z) \bar{g}(z) d\vartheta(z) d\sigma(\zeta) \\
&= \int_{\mathbb{D}^n} f(z) \bar{g}(z) d\vartheta(z).
\end{aligned}$$

This shows that $\langle f - f_s, g \rangle = 0$ for all $g \in \mathcal{H}_s$. Since f_s belongs to $\mathcal{H}_s$, we conclude that $f_s = Q_s f$.

Next, suppose $l \neq k$. Let $f \in \mathcal{H}_l$ and $g \in \mathcal{H}_k$. For any $\zeta \in \mathbb{T}^n$, we have

$$\begin{aligned}
\int_{\mathbb{D}^n} \zeta^{l-k} f(z) \bar{g}(z) d\vartheta(z) &= \int_{\mathbb{D}^n} f(z \cdot \zeta) \bar{g}(z \cdot \zeta) d\vartheta(z) \\
&= \int_{\mathbb{D}^n} f(z) \bar{g}(z) d\vartheta(z).
\end{aligned}$$

Since $l \neq k$, we conclude that $\int_{\mathbb{D}^n} f(z) \bar{g}(z) d\vartheta(z) = 0$. Thus, $\mathcal{H}_l \perp \mathcal{H}_k$.

To show $L_\vartheta^2 = \bigoplus_{l \in \mathbb{Z}^n} \mathcal{H}_l$, it suffices to show that for any $f \in L_\vartheta^2$, the identity $\|f\|_2^2 = \sum_{l \in \mathbb{Z}^n} \|Q_l(f)\|_2^2$ holds true. Indeed, for $f \in L_\vartheta^2$,

$$\begin{aligned}
\|f\|_2^2 &= \int_{\mathbb{D}^n} |f(z)|^2 d\vartheta(z) \\
&= \int_{\mathbb{D}^n} \int_{\mathbb{T}^n} |f(z \cdot \zeta)|^2 d\sigma(\zeta) d\vartheta(z) \\
&= \int_{\mathbb{D}^n} \sum_{l \in \mathbb{Z}^n} \left| \int_{\mathbb{T}^n} f(z \cdot \zeta) \bar{\zeta}^l d\sigma(\zeta) \right|^2 d\vartheta(z) \\
&\quad (\text{since } \{\zeta^l : l \in \mathbb{Z}^n\} \text{ is an orthonormal basis for } L^2(\mathbb{T}^n, \sigma)) \\
&= \sum_{l \in \mathbb{Z}^n} \int_{\mathbb{D}^n} \left| \int_{\mathbb{T}^n} f(z \cdot \zeta) \bar{\zeta}^l d\sigma(\zeta) \right|^2 d\vartheta(z) \\
&= \sum_{l \in \mathbb{Z}^n} \|Q_l(f)\|_2^2. \quad \square
\end{aligned}$$

It follows from the proof of Lemma 2.2 that for each $s \in \mathbb{Z}^n$, there is a function f_s such that $f_s(z \cdot \gamma) = \gamma^s f_s(z)$ for all $z \in \mathbb{D}^n$ and all $\zeta \in \mathbb{T}^n$ and $Q_s(f)(z) = f_s(z)$ for ϑ -a.e. z . If f is continuous on the closed polydisc $\overline{\mathbb{D}^n}$, then the integral in (2.2) is well-defined for all z in $\overline{\mathbb{D}^n}$ and f_s is also

continuous on $\overline{\mathbb{D}}^n$. We have seen that the series $\sum_{s \in \mathbb{Z}^n} f_s$ converges to f in the $L^2_{\mathcal{H}}$ -norm. In general, for f in $C(\overline{\mathbb{D}}^n)$, the series does not converge uniformly to f . However, the Cesàro means of the functions $\{f_s : s \in \mathbb{Z}^n\}$ do converge uniformly to f as we will see next.

For any integer $N \geq 1$, the Cesàro mean $\Lambda_N(f)$ is defined by the formula

$$\begin{aligned} \Lambda_N(f)(z) &= \sum_{|s_1|, \dots, |s_n| \leq N} \left(1 - \frac{|s_1|}{N+1}\right) \cdots \left(1 - \frac{|s_n|}{N+1}\right) f_{s_1, \dots, s_n}(z) \\ &= \int_{\mathbb{T}^n} \left\{ \sum_{|s_1|, \dots, |s_n| \leq N} \left(1 - \frac{|s_1|}{N+1}\right) \cdots \left(1 - \frac{|s_n|}{N+1}\right) \zeta_1^{s_1} \cdots \zeta_n^{s_n} \right\} f(z \cdot \zeta) d\sigma(\zeta) \\ &= \int_{\mathbb{T}^n} \mathbf{F}_N(\zeta_1) \cdots \mathbf{F}_N(\zeta_n) f(z \cdot \zeta) d\sigma(\zeta), \end{aligned}$$

where $\mathbf{F}_N$ is the N th Fejér's kernel. It now follows from a well known result in harmonic analysis (see, for example, Sections 2.2 and 9.2 in [5]) that $\Lambda_N(f) \rightarrow f$ uniformly on $\overline{\mathbb{D}}^n$ as $N \rightarrow \infty$ if f is continuous on $\overline{\mathbb{D}}^n$.

3. HANKEL OPERATORS WITH QUASI-HOMOGENEOUS SYMBOLS

Recall from Section 2 that $A_{\mathcal{H}}^2$ has the standard orthonormal basis consisting of monomials $\{e_m(z) = \frac{z^m}{\sqrt{c_m}} : m \in \mathbb{Z}_+^n\}$, where

$$c_m = \int_{[0,1]^n} r_1^{2m_1} \cdots r_n^{2m_n} d\mu(r_1, \dots, r_n).$$

We also recall that for $l \in \mathbb{Z}^n$, Q_l is the orthogonal projection from $L^2_{\mathcal{H}}$ onto the subspace $\mathcal{H}_l$ of quasi-homogeneous functions of multi-degree l .

For two n -tuples of integers $l = (l_1, \dots, l_n)$ and $s = (s_1, \dots, s_n)$, we write $l \succeq s$ if $l_j \geq s_j$ for all $1 \leq j \leq n$ and $l \not\succeq s$ if otherwise. We will also use 0 to denote $(0, \dots, 0)$. For $m \in \mathbb{Z}_+^n$ and $l \in \mathbb{Z}^n$, $Q_l(e_m)$ is either 0 (when $l \neq m$) or e_m (when $l = m$). Thus, $Q_l(A_{\mathcal{H}}^2) = \{0\}$ if $l \not\succeq 0$ and $Q_l(A_{\mathcal{H}}^2) = \mathbb{C}e_l$ if $l \succeq 0$. This shows that $A_{\mathcal{H}}^2$ is an invariant subspace for Q_l , hence it is also a reducing subspace since Q_l is a projection. Let P be the orthogonal projection from L^2 onto $A_{\mathcal{H}}^2$. Then we have $PQ_l = Q_lP$ and this in turn shows that $\mathcal{H}_l$ is a reducing subspace for P .

Lemma 3.1. *Let s be in $\mathbb{Z}^n$. Suppose f is a bounded function on $\mathbb{D}^n$ so that we have $f(r_1\zeta_1, \dots, r_n\zeta_n) = \zeta^s f(r_1, \dots, r_n)$ for all $z = (r_1\zeta_1, \dots, r_n\zeta_n)$ in $\mathbb{D}^n$. Then $H_f^* H_f$ is a diagonal operator with respect to the standard orthonormal basis. The eigenvalues of $H_f^* H_f$ are given by*

$$\lambda_m = \frac{1}{c_m} \int_{[0,1]^n} |f(t_1, \dots, t_n)|^2 t_1^{2m_1} \cdots t_n^{2m_n} d\mu(t_1, \dots, t_n)$$

if $m + s \not\geq 0$ and

$$\lambda_m = \frac{1}{c_m} \int_{[0,1]^n} |f(t_1, \dots, t_n)|^2 t_1^{2m_1} \dots t_n^{2m_n} d\mu(t_1, \dots, t_n) \\ - \frac{1}{c_m c_{m+s}} \left| \int_{[0,1]^n} f(t_1, \dots, t_n) t_1^{2m_1+s_1} \dots t_n^{2m_n+s_n} d\mu(t_1, \dots, t_n) \right|^2$$

if $m + s \geq 0$.

Proof. For any $m \in \mathbb{Z}_+^n$, $f e_m$ belongs to $\mathcal{H}_{s+m}$, which is an invariant subspace for P . Therefore, $P(f e_m)$ and $H_f e_m = f e_m - P(f e_m)$ also belong to $\mathcal{H}_{s+m}$. We have

$$P(f e_m) = \sum_{k \in \mathbb{Z}_+^n} \langle P(f e_m), e_k \rangle e_k = \begin{cases} 0 & \text{if } s + m \not\geq 0, \\ \langle f e_m, e_{s+m} \rangle e_{s+m} & \text{if } s + m \geq 0. \end{cases}$$

Now for $k \neq m$, $\langle H_f^* H_f e_m, e_k \rangle = \langle H_f e_m, H_f e_k \rangle = 0$ since $H_f e_m \in \mathcal{H}_{m+s}$, $H_f e_k \in \mathcal{H}_{k+s}$ and $\mathcal{H}_{m+s} \perp \mathcal{H}_{k+s}$ by Lemma 2.2. Thus, $H_f^* H_f$ is a diagonal operator with respect to the standard orthonormal basis $\{e_m : m \in \mathbb{Z}_+^n\}$. The eigenvalues of $H_f^* H_f$ are given by

$$\lambda_m = \langle H_f^* H_f e_m, e_m \rangle = \|H_f e_m\|_2^2 \\ = \|f e_m\|_2^2 - \|P(f e_m)\|_2^2 \\ = \begin{cases} \|f e_m\|_2^2 & \text{if } s + m \not\geq 0, \\ \|f e_m\|_2^2 - |\langle f e_m, e_{s+m} \rangle|^2 & \text{if } s + m \geq 0, \end{cases}$$

for $m \in \mathbb{Z}_+^n$. Since

$$\|f e_m\|_2^2 = \frac{1}{c_m} \int_{[0,1]^n} |f(t_1, \dots, t_n)|^2 t_1^{2m_1} \dots t_n^{2m_n} d\mu(t_1, \dots, t_n),$$

and

$$\langle f e_m, e_{s+m} \rangle = \frac{1}{\sqrt{c_m c_{s+m}}} \int_{[0,1]^n} f(t_1, \dots, t_n) t_1^{2m_1+s_1} \dots t_n^{2m_n+s_n} d\mu(t_1, \dots, t_n),$$

the conclusion of the lemma follows. $\square$

Remark 3.2. Let us consider the case $n = 1$ and $f(z) = z^u \bar{z}^v$ for integers $u, v \geq 0$. We see that f belongs to $\mathcal{H}_s$ with $s = u - v$. From Lemma 3.1, $H_f^* H_f$ is a diagonal operator with eigenvalues λ_m for $m \in \mathbb{Z}_+$. For all positive integers $m \geq v - u$, we have

$$\lambda_m = \frac{\int_{[0,1]} t^{2m+2(u+v)} d\mu(t)}{\int_{[0,1]} t^{2m} d\mu(t)} - \frac{\left| \int_{[0,1]} t^{2m+2u} d\mu(t) \right|^2}{\left(\int_{[0,1]} t^{2m} d\mu(t) \right) \left(\int_{[0,1]} t^{2m+2(u-v)} d\mu(t) \right)}.$$

Since $\mu([r, 1]) > 0$ for all $0 < r < 1$, it can be showed that $\lim_{m \rightarrow \infty} \lambda_m = 0$ (see Lemma 3.3 below). Therefore, $H_f^* H_f$ is a compact operator, which implies that H_f is also a compact operator. Thus, for any polynomial

$p = p(z, \bar{z})$, H_p is compact. Since any function g in $C(\overline{\mathbb{D}^n})$ can be uniformly approximated by polynomials, we conclude that H_g is also a compact operator.

Our characterization of compactness of Hankel operators when $n \geq 2$ depends on the following lemma. For a sketch of its proof when $\delta = \beta = (0, \dots, 0)$, see Lemma 2.4 in [6]. The proof for arbitrary δ, β is similar.

Lemma 3.3. *Let $\mu_1, \dots, \mu_N$ be positive measures on $[0, 1)$ so that $\mu_j([r, 1)) > 0$ for all $0 < r < 1$, all $1 \leq j \leq N$. Suppose φ is a function on $[0, 1)^N$ so that*

$$\lim_{(r_1, \dots, r_N) \rightarrow (1, \dots, 1)} \varphi(r_1, \dots, r_N) = \alpha. \text{ Then for any } N\text{-tuples of real numbers}$$

$\delta = (\delta_1, \dots, \delta_N)$ and $\beta = (\beta_1, \dots, \beta_N)$, we have

$$\lim_{(m_1, \dots, m_N) \rightarrow (\infty, \dots, \infty)} \frac{\int_{[0, 1)^N} \varphi(r) r_1^{m_1 + \delta_1} \dots r_N^{m_N + \delta_N} d\mu(r)}{\int_{[0, 1)^N} r_1^{m_1 + \beta_1} \dots r_N^{m_N + \beta_N} d\mu(r)} = \alpha.$$

In the rest of the paper, we will consider only measures μ of the form $d\mu(r_1, \dots, r_n) = d\mu_1(r_1) \dots d\mu_n(r_n)$, where $\mu_1, \dots, \mu_n$ are regular Borel probability measures on the interval $[0, 1)$ such that $\mu_j([r, 1)) > 0$ for all $0 < r < 1$ and $1 \leq j \leq n$. Recall that ϑ is the measure on $\mathbb{D}^n$ that is related to μ by equation (1.1). We now define a measure γ on the topological boundary $\partial\mathbb{D}^n$ associated with ϑ . It is clear that $\partial\mathbb{D}^n$ is the disjoint union of $2^n - 1$ parts of the form $A_1 \times \dots \times A_n$, where A_j is either $\mathbb{T}$ or $\mathbb{D}$ and not all are $\mathbb{D}$. Suppose $W = A_1 \times \dots \times A_n$ is a part of $\partial\mathbb{D}^n$. Let $1 \leq j_1 < \dots < j_p \leq n$ be integers so that $A_{j_1} = \dots = A_{j_p} = \mathbb{D}$ and $A_j = \mathbb{T}$ if $j \notin \{j_1, \dots, j_p\}$. We define γ to be the unique regular Borel measure on W that satisfies

$$\begin{aligned} & \int_W f(w) d\gamma(w) \\ &= \int_{[0, 1)^p} \left\{ \int_{\mathbb{T}^n} f(\zeta_1, \dots, r_{j_1} \zeta_{j_1}, \dots, r_{j_p} \zeta_{j_p}, \dots, \zeta_n) d\sigma(\zeta) \right\} d\mu_{j_1}(r_{j_1}) \dots d\mu_{j_p}(r_{j_p}) \end{aligned}$$

for all $f \in C_c(W)$.

The following theorem characterizes compact Hankel operators with continuous quasi-homogeneous symbols when $n \geq 2$.

Theorem 3.4. *Suppose $n \geq 2$ and $s \in \mathbb{Z}^n$. Suppose $f \in \mathcal{H}_s$ is continuous on $\overline{\mathbb{D}^n}$ such that H_f is compact. If $s \succeq 0$, then $f(z) = f(1, \dots, 1)z^s$ for γ -a.e. z in $\partial\mathbb{D}^n$. If $s \not\succeq 0$, then $f(z) = 0$ for γ -a.e. z in $\partial\mathbb{D}^n$.*

Proof. By the remark after the proof of Lemma 2.2, we may assume that $f(\zeta_1 z_1, \dots, \zeta_n z_n) = \zeta^s f(z_1, \dots, z_n)$ for all $\zeta \in \mathbb{T}^n$ and all $z \in \overline{\mathbb{D}^n}$. In particular, for all $z = (r_1 \zeta_1, \dots, r_n \zeta_n)$ in $\overline{\mathbb{D}^n}$, we have

$$f(r_1 \zeta_1, \dots, r_n \zeta_n) = \zeta^s f(r_1, \dots, r_n). \quad (3.1)$$

Suppose $1 \leq j < n$. Put $m_1 = \max\{0, -s_1\}, \dots, m_j = \max\{0, -s_j\}$. Since H_f is compact, $H_f^* H_f$ is also compact. By Lemma 3.1, $H_f^* H_f$ is diagonalizable and its eigenvalues are λ_m 's for $m = (m_1, \dots, m_j, m_{j+1}, \dots, m_n)$.

Therefore, $\lim_{(m_{j+1}, \dots, m_n) \rightarrow (\infty, \dots, \infty)} \lambda_m = 0$. Using the formula for λ_m (when $m + s \succeq 0$) in Lemma 3.1 together with Lemma 3.3 and the fact that $c_m = \int_{[0,1]^n} t_1^{2m_1} \cdots t_n^{2m_n} d\mu_1(t_1) \cdots d\mu_n(t_n)$, we conclude that

$$\begin{aligned} & \int_{[0,1]^j} |f(t_1, \dots, t_j, 1, \dots, 1)|^2 t_1^{2m_1} \cdots t_j^{2m_j} d\mu_1(t_1) \cdots d\mu_j(t_j) \\ &= \frac{\left| \int_{[0,1]^j} f(t_1, \dots, t_j, 1, \dots, 1) t_1^{2m_1+s_1} \cdots t_j^{2m_j+s_j} d\mu_1(t_1) \cdots d\mu_j(t_j) \right|^2}{\int_{[0,1]^j} t_1^{2m_1+2s_1} \cdots t_j^{2m_j+2s_j} d\mu_1(t_1) \cdots d\mu_j(t_j)}. \end{aligned}$$

Let $F(t) = f(t_1, \dots, t_j, 1, \dots, 1) t_1^{m_1} \cdots t_j^{m_j}$ and $G(t) = t_1^{m_1+s_1} \cdots t_j^{m_j+s_j}$ for $t = (t_1, \dots, t_j) \in [0,1]^j$. Then the above identity shows that

$$\begin{aligned} & \left\{ \int_{[0,1]^j} |F(t)|^2 d\mu_1(t_1) \cdots d\mu_j(t_j) \right\} \left\{ \int_{[0,1]^n} G^2(t) d\mu_1(t_1) \cdots d\mu_j(t_j) \right\} \\ &= \left| \int_{[0,1]^j} (FG)(t) d\mu_1(t_1) \cdots d\mu_j(t_j) \right|^2. \end{aligned}$$

This means that Holder's inequality applied to F and G is in fact an equality. Therefore, we have $F(t) = \alpha G(t)$, or equivalently,

$$f(t_1, \dots, t_j, 1, \dots, 1) t_1^{m_1} \cdots t_j^{m_j} = \alpha t_1^{m_1+s_1} \cdots t_j^{m_j+s_j} \quad (3.2)$$

for $\mu_1 \times \cdots \times \mu_j$ -a.e. $t = (t_1, \dots, t_j) \in [0,1]^j$, where α is a constant. Since 1 belongs to the support of all the measures $\mu_1, \dots, \mu_j$, we may let $t_1 = \cdots = t_j = 1$ to obtain $\alpha = f(1, \dots, 1)$. Recall that in (3.2), $m_1 = \max\{0, -s_1\}, \dots, m_j = \max\{0, -s_j\}$.

Suppose first $s_1, \dots, s_n \geq 0$. Since $m_1 = \cdots = m_j = 0$, we obtain from (3.2) that $f(t_1, \dots, t_j, 1, \dots, 1) = f(1, \dots, 1) t_1^{s_1} \cdots t_j^{s_j}$ for $\mu_1 \times \cdots \times \mu_j$ -a.e. $t = (t_1, \dots, t_j) \in [0,1]^j$. This together with (3.1) implies that $f(z) = f(1, \dots, 1) z^s$ for γ -a.e. z in $W = \mathbb{D}^j \times \mathbb{T}^{n-j}$, which is a part of the boundary $\partial \mathbb{D}^n$.

Now suppose $s_p < 0$ for some $1 \leq p \leq n$. We will show that in this case $f(1, \dots, 1) = 0$. Without loss of generality, we may assume $p = 1$. For all large positive integers $m_2, \dots, m_n$, let $m = (0, m_2, \dots, m_n)$ (the assumption that $n \geq 2$ is needed here). Since $m + s \not\succeq 0$,

$$\lambda_m = \frac{\int_{[0,1]^n} |f(t_1, \dots, t_n)|^2 t_2^{2m_2} \cdots t_n^{2m_n} d\mu_1(t_1) \cdots d\mu_n(t_n)}{\int_{[0,1]^n} t_2^{2m_2} \cdots t_n^{2m_n} d\mu_1(t_1) \cdots d\mu_n(t_n)}.$$

Letting $(m_2, \dots, m_n) \rightarrow (\infty, \dots, \infty)$ and using Lemma 3.3 together with the fact that $\lambda_m \rightarrow 0$, we conclude that

$$0 = \int_{[0,1]} |f(t_1, 1, \dots, 1)|^2 d\mu_1(t_1).$$

This implies $f(t_1, 1, \dots, 1) = 0$ for μ_1 -a.e. t_1 on $[0, 1]$. Since 1 is in the support of μ_1 and f is continuous at the point $(1, \dots, 1)$, it follows that $f(1, \dots, 1) = 0$.

Now (3.2) gives $f(t_1, \dots, t_j, 1, \dots, 1)t_1^{m_1} \dots t_j^{m_j} = 0$ for $\mu_1 \times \dots \times \mu_j$ -a.e. $t = (t_1, \dots, t_j)$ in $[0, 1]^j$. For such t , if $t_1^{m_1} \dots t_j^{m_j} \neq 0$, then we have $f(t_1, \dots, t_j, 1, \dots, 1) = 0$. Otherwise, there exists $1 \leq i \leq j$ so that $t_i = 0$ and $m_i > 0$. But $m_j = \max\{0, -s_i\}$, so $s_i < 0$. Since $t_i = 0 = t_i \zeta_i$ for any $|\zeta_i| = 1$, we have

$$\begin{aligned} f(t_1, \dots, t_i, \dots, t_j, 1, \dots, 1) &= f(t_1, \dots, t_i \zeta_i, \dots, t_j, 1, \dots, 1) \\ &= \zeta_i^{s_i} f(t_1, \dots, t_i, \dots, t_j, 1, \dots, 1). \end{aligned}$$

This implies $f(t_1, \dots, t_j, 1, \dots, 1) = 0$ because ζ_i can be chosen so that $\zeta_i^{s_i} \neq 1$. Therefore, $f(t_1, \dots, t_j, 1, \dots, 1) = 0$ for $\mu_1 \times \dots \times \mu_j$ -a.e. $t = (t_1, \dots, t_j)$ in $[0, 1]^j$, which implies $f(z) = 0$ for γ -a.e. $z \in W = \mathbb{D}^j \times \mathbb{T}^{n-j}$.

The same argument applies to other parts of $\partial \mathbb{D}^n$ which are different from $\mathbb{T}^n$. On $\mathbb{T}^n$, (3.1) gives $f(\zeta) = \zeta^s f(1, \dots, 1)$. If $s_p < 0$ for some $1 \leq p \leq n$ then since $f(1, \dots, 1) = 0$ as showed above, we conclude that $f(\zeta) = 0$ for $\zeta \in \mathbb{T}^n$. So the conclusions of the proposition also hold for z in $\mathbb{T}^n \subset \partial \mathbb{D}^n$. The proof of the proposition is now completed. $\square$

4. COMPACT HANKEL OPERATORS WITH MORE GENERAL SYMBOLS

We have seen that any f in $L_{\mathcal{Q}}^2$ admits the decomposition $f = \sum_{l \in \mathbb{Z}^n} f_l$, where $f_l = Q_l(f)$ is the orthogonal projection of f on the space of quasi-homogeneous functions of multi-degree l . The next proposition shows that the compactness of H_f implies the compactness of each H_{f_l} . We are then able to apply the results in the previous section. The Hardy space version of the proposition was proved in [1]. Our proof here, which also works for the Hardy space, is more direct.

Proposition 4.1. *Suppose $f \in L_{\mathcal{Q}}^2$ so that the operator H_f is compact on $A_{\mathcal{Q}}^2$. Then for each $s \in \mathbb{Z}^n$, the operator H_{f_s} is compact.*

Proof. For any $m \in \mathbb{Z}_+^n$, we have

$$\begin{aligned} H_f e_m &= (I - P)(f e_m) = (I - P)\left(\sum_{l \in \mathbb{Z}^n} f_l\right) e_m \\ &= \sum_{l \in \mathbb{Z}^n} (I - P)(f_l e_m) = \sum_{l \in \mathbb{Z}^n} H_{f_l} e_m. \end{aligned}$$

Since $Q_{s+m}(H_{f_l} e_m) = 0$ if $l \neq s$ and $Q_{s+m}(H_{f_s} e_m) = H_{f_s} e_m$, we obtain $Q_{s+m}(H_f e_m) = H_{f_s} e_m$. From Lemma 3.1, $H_{f_s}^* H_{f_s}$ is a diagonal operator with eigenvalues λ_m given by

$$\lambda_m = \|H_{f_s} e_m\|_2^2 = \|Q_{s+m}(H_f e_m)\|_2^2 \leq \|H_f e_m\|_2^2.$$

Since H_f is compact, we have $\lim_{|m| \rightarrow \infty} \|H_f e_m\|_2 = 0$ (here $|m| = m_1 + \dots + m_n$). This implies $\lim_{|m| \rightarrow \infty} \lambda_m = 0$ and hence, $H_{f_s}^* H_{f_s}$ is compact. Thus, H_{f_s} is a compact operator. $\square$

Suppose g belongs to A_{ϑ}^2 . It was showed by Axler [2] that when $n = 1$ and ϑ is the Lebesgue measure on the disk $\mathbb{D}$, $H_{\bar{g}}$ is compact if and only if g is in the little Bloch space, that is, $\lim_{|z| \uparrow 1} (1 - |z|^2)g'(z) = 0$. For $n \geq 2$ and ϑ is the Lebesgue measure on the polydisc $\mathbb{D}^n$, a special case of [3, Theorem D] gives that $H_{\bar{g}}$ is compact if and only if g is a constant function and in this case, $H_{\bar{g}} = 0$. The following corollary to Proposition 4.1 shows that this holds true for general measures ϑ .

Corollary 4.2. *Suppose $n \geq 2$. Let g be a function in A_{ϑ}^2 so that $H_{\bar{g}}$ is compact. Then g is a constant function and hence $H_{\bar{g}} = 0$.*

Proof. Write $g = \sum_{m \in \mathbb{Z}_+^n} c_m e_m$. For $m \in \mathbb{Z}_+^n$, since $Q_{-m}(\bar{g}) = \bar{c}_m \bar{e}_m$ and $H_{\bar{g}}$ is compact, Proposition 4.1 implies that $H_{\bar{c}_m \bar{e}_m}$ is compact. Theorem 3.4 then shows that $\bar{c}_m = 0$ for all $m \neq 0$. Thus, g is a constant function. $\square$

We are now ready for our main theorem in this paper.

Theorem 4.3. *Suppose $n \geq 2$. Let f be continuous on $\overline{\mathbb{D}^n}$ so that H_f is a compact operator. Then there is a function h which is continuous on $\overline{\mathbb{D}^n}$ and holomorphic on $\mathbb{D}^n$, and a bounded function g satisfying $\lim_{z \rightarrow \partial \mathbb{D}^n} g(z) = 0$ so that $f(z) = h(z) + g(z)$ for ϑ -a.e. z in $\mathbb{D}^n$.*

Proof. For any $s \in \mathbb{Z}^n$, Proposition 4.1 shows that H_{f_s} is compact. Since f is continuous, each f_s is also continuous. By Theorem 3.4, there is a holomorphic monomial h_s so that $(f_s - h_s)(w) = 0$ for γ -a.e. $w \in \partial \mathbb{D}^n$. (In fact, $h_s = 0$ if $s \not\geq 0$ and $h_s(w) = f_s(1, \dots, 1)w^s$ if $s \geq 0$).

For each integer $N \geq 1$, define

$$p_N(z) = \sum_{|s_1|, \dots, |s_n| \leq N} \left(1 - \frac{|s_1|}{N+1}\right) \cdots \left(1 - \frac{|s_n|}{N+1}\right) h_{s_1, \dots, s_n}(z) \quad \text{for } z \in \overline{\mathbb{D}^n}.$$

Then p_N is a holomorphic polynomial and $p_N(w) = \Lambda_N(f)(w)$ for γ -a.e. $w \in \partial \mathbb{D}^n$, where $\Lambda_N(f)$ is the N th Cesàro mean of f . Since γ restricted on $\mathbb{T}^n \subset \partial \mathbb{D}^n$ is the surface measure and $p_N - \Lambda_N(f)$ is continuous, we actually have $p_N(w) = \Lambda_N(f)(w)$ for all $w \in \mathbb{T}^n$. By the remark at the end of Section 2, $\Lambda_N(f)$ converges to f uniformly on $\overline{\mathbb{D}^n}$. In particular, $p_N|_{\mathbb{T}^n} = \Lambda_N(f)|_{\mathbb{T}^n}$ converges to $f|_{\mathbb{T}^n}$ uniformly. This implies that there is a function h which is continuous on $\overline{\mathbb{D}^n}$ and holomorphic on $\mathbb{D}^n$ so that p_N converges uniformly to h on $\overline{\mathbb{D}^n}$. Since $p_N(w) = \Lambda_N(f)(w)$ for γ -a.e. $w \in \partial \mathbb{D}^n$ for each N , we conclude that $h(w) = f(w)$ for γ -a.e. $w \in \partial \mathbb{D}^n$. Let $\tilde{g} = f - h$. Then $\tilde{g}$ is continuous on $\overline{\mathbb{D}^n}$ and $\tilde{g}(w) = 0$ for γ -a.e. w on $\partial \mathbb{D}^n$. By Lemma 2.1 in [6], there is a function g such that $\tilde{g}(z) = g(z)$ for ϑ -a.e. z in $\mathbb{D}^n$ and $\lim_{z \rightarrow w} g(z) = 0$ for all $w \in \partial \mathbb{D}^n$. From the compactness of $\partial \mathbb{D}^n$, it follows that $\lim_{z \rightarrow \partial \mathbb{D}^n} g(z) = 0$. We then have $f(z) = h(z) + \tilde{g}(z) = h(z) + g(z)$

for ϑ -a.e. $z \in \mathbb{D}^n$ and h, g satisfy the requirements in the conclusion of the theorem. $\square$

The continuity of f on $\overline{\mathbb{D}^n}$ in Theorem 4.3 cannot be dropped. Indeed, there are bounded functions f which are continuous on the open polydisc $\mathbb{D}^n$ such that H_f is compact and no decomposition $f = h + g$ with $h \in A_{\vartheta}^2$ and $\lim_{z \rightarrow \partial \mathbb{D}^n} g(z) = 0$ is possible. In the rest of the section, we will give a construction of such a function.

Let $0 < r_1 < r_2 < \dots$ be an increasing sequence of positive real numbers that converges to 1. Set $j_1 = 1$. Since $(r_{j_1}, 1)^n = \cup_{j=j_1+1}^{\infty} (r_{j_1}, r_j)^n$ and $\mu((r_{j_1}, 1)^n) > 0$, there is an integer $j_2 \geq j_1 + 1$ so that $\mu((r_{j_1}, r_{j_2})^n) > 0$. Since $(r_{j_2}, 1)^n = \cup_{j=j_2+1}^{\infty} (r_{j_2}, r_j)^n$ and $\mu((r_{j_2}, 1)^n) > 0$, there is an integer $j_3 \geq j_2 + 1$ so that $\mu((r_{j_2}, r_{j_3})^n) > 0$. Continuing this process, we find a sequence of integers $\{j_k\}_{k=1}^{\infty}$ such that $j_{k+1} \geq j_k + 1$ and $\mu((r_{j_k}, r_{j_{k+1}})^n) > 0$ for $k = 1, 2, \dots$. For each such k , let $R_k = (r_{j_k}, r_{j_{k+1}})^n$. Choose an open subset V_k of $\mathbb{T}^n$ so that $0 < \sigma(V_k) < 1/k$ and $\int_{E_k} K_z(z) d\vartheta(z) < 1/k^2$, where $E_k = \{r \cdot \zeta : r \in R_k, \zeta \in V_k\}$. The existence of V_k follows from the fact that the function $z \mapsto K_z(z)$ is bounded on compact sets. Since E_k is open in $\mathbb{D}^n$ and $\vartheta(E_k) > 0$, using the regularity of ϑ , we can choose a continuous function $0 \leq f_k \leq 1$ so that f_k is supported in E_k and $\vartheta(\{z : f(z) = 1\}) > 0$. Put $f = \sum_{k=1}^{\infty} f_k$. Since the sets E_k 's are pairwise disjoint, the function f is continuous on $\mathbb{D}^n$ and $0 \leq f(z) \leq 1$ for all $z \in \mathbb{D}^n$.

We now show that H_f is a Hilbert-Schmidt operator, hence it is compact. Indeed, from (2.1),

$$\begin{aligned} \sum_{m \in \mathbb{Z}_+^n} \|H_f e_m\|_2^2 &\leq \int_{\mathbb{D}^n} |f(z)|^2 K_z(z) d\vartheta(z) \\ &\leq \sum_{k=1}^{\infty} \int_{E_k} |f(z)|^2 K_z(z) d\vartheta(z) < \sum_{k=1}^{\infty} \frac{1}{k^2} < \infty. \end{aligned}$$

For each $s \in \mathbb{Z}^n$, we will show that $\lim_{z \rightarrow \partial \mathbb{D}^n} Q_s(f)(z) = 0$. Since $f \geq 0$, it follows from formula (2.2) in Lemma 2.2 that $|Q_s(f)(z)| \leq |Q_0(f)(z)|$. So it suffices to prove $\lim_{z \rightarrow \partial \mathbb{D}^n} Q_0(f)(z) = 0$. For $z = (z_1, \dots, z_n) \in \mathbb{D}^n$, we have

$$\begin{aligned} Q_0(f)(z) &= \int_{\mathbb{T}^n} f(z \cdot \zeta) d\sigma(\zeta) = \int_{\mathbb{T}^n} f(|z_1|\zeta_1, \dots, |z_n|\zeta_n) d\sigma(\zeta) \\ &\leq \sum_{k=1}^{\infty} \int_{\mathbb{T}^n} \chi_{E_k}(|z_1|\zeta_1, \dots, |z_n|\zeta_n) d\sigma(\zeta). \end{aligned}$$

By the definition of E_k , $\chi_{E_k}(|z_1|\zeta_1, \dots, |z_n|\zeta_n) = 1$ if and only if the n -tuple $(|z_1|, \dots, |z_n|)$ belongs to $R_k = (r_{j_k}, r_{j_{k+1}})^n$ and $(\zeta_1, \dots, \zeta_n)$ belongs to V_k . Let $k_0 \geq 2$ be a positive integer. Suppose $z = (z_1, \dots, z_n)$ so that $|z_i| > r_{j_{k_0}}$ for some $1 \leq i \leq n$. If $(|z_1|, \dots, |z_n|)$ does not belong to any R_k , $k \geq 1$, then $Q_0(f)(z) = 0$. Otherwise, there is exactly one k so that $(|z_1|, \dots, |z_n|) \in R_k$,

which is $(r_{j_k}, r_{j_{k+1}})^n$. Since $|z_i| > r_{j_{k_0}}$, we conclude that $r_{j_{k+1}} > r_{k_0}$, which implies $k \geq k_0$. Therefore,

$$|Q_0(f)(z)| \leq \int_{\mathbb{T}^n} \chi_{E_k}(|z_1|\zeta_1, \dots, |z_n|\zeta_n) d\sigma(\zeta) = \sigma(V_k) \leq \frac{1}{k} \leq \frac{1}{k_0}.$$

Since this holds true for any z that does not belong to the compact set $[0, r_{j_{k_0}}]^n \times \mathbb{T}^n$, we conclude that $\lim_{z \rightarrow \partial \mathbb{D}^n} Q_0(f)(z) = 0$.

Suppose there were a decomposition $f = h + g$, where $h \in A_{\vartheta}^2$ and $g(z) \rightarrow 0$ as $z \rightarrow \partial \mathbb{D}^n$. We will show that there would be a contraction. For $s \succeq 0$, from formula (2.2) in Lemma 2.2 we see that $Q_s(g)(z) \rightarrow 0$ as $z \rightarrow \partial \mathbb{D}^n$. This implies $Q_s(h)(z) = Q_s(f)(z) - Q_s(g)(z) \rightarrow 0$ as $z \rightarrow \partial \mathbb{D}^n$. But $Q_s(h)$ is a multiple of z^s , as explained at the beginning of Section 3, so we have $Q_s(h) = 0$ for all $s \succeq 0$. It follows that $h = 0$ and hence, $f = g$. This is a contradiction because $g(z) \rightarrow 0$ as $z \rightarrow \partial \mathbb{D}^n$ but by the construction of f , for any compact subset $M \subset \mathbb{D}^n$, the set $\{z \in \mathbb{D}^n \setminus M : f(z) = 1\}$ has positive ϑ -measure.

REFERENCES

- [1] Patrick Ahern, El Hassan Youssfi, and Kehe Zhu, *Compactness of Hankel operators on Hardy-Sobolev spaces of the polydisk*, J. Operator Theory **61** (2009), no. 2, 301–312.
- [2] Sheldon Axler, *The Bergman space, the Bloch space, and commutators of multiplication operators*, Duke Math. J. **53** (1986), no. 2, 315–332. MR 850538 (87m:47064)
- [3] D. Békollé, C. A. Berger, L. A. Coburn, and K. H. Zhu, *BMO in the Bergman metric on bounded symmetric domains*, J. Funct. Anal. **93** (1990), no. 2, 310–350. MR 1073289 (91j:32034)
- [4] Mischa Cotlar and Cora Sadosky, *Abstract, weighted, and multidimensional Adamjan-Arov-Kreĭn theorems, and the singular numbers of Sarason commutants*, Integral Equations Operator Theory **17** (1993), no. 2, 169–201. MR 1233667 (94k:47040)
- [5] Yitzhak Katznelson, *An introduction to harmonic analysis*, third ed., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 2004. MR 2039503 (2005d:43001)
- [6] Trieu Le, *On Toeplitz operators on Bergman spaces of the unit polydisk*, Proc. Amer. Math. Soc. **138** (2010), 275–285.
- [7] Issam Louhichi, Elizabeth Strouse, and Lova Zakariasy, *Products of Toeplitz operators on the Bergman space*, Integral Equations Operator Theory **54** (2006), no. 4, 525–539. MR 2222982 (2007a:47033)
- [8] Karel Stroethoff, *Compact Hankel operators on the Bergman space*, Illinois J. Math. **34** (1990), no. 1, 159–174. MR 1031892 (91a:47030)
- [9] ———, *Compact Hankel operators on the Bergman spaces of the unit ball and polydisk in $\mathbf{C}^n$* , J. Operator Theory **23** (1990), no. 1, 153–170. MR 1054822 (91i:47040)
- [10] De Chao Zheng, *Toeplitz operators and Hankel operators*, Integral Equations Operator Theory **12** (1989), no. 2, 280–299. MR 986598 (90b:47044)
- [11] Kehe Zhu, *Operator theory in function spaces*, second ed., Mathematical Surveys and Monographs, vol. 138, American Mathematical Society, Providence, RI, 2007. MR 2311536

DEPARTMENT OF MATHEMATICS, MAIL STOP 942, UNIVERSITY OF TOLEDO, TOLEDO, OH 43606

E-mail address: `trieu.le2@utoledo.edu`